

Positive toric fibrations

Misha Verbitsky¹

verbit@maths.gla.ac.uk, verbit@mccme.ru

Abstract

A principal toric bundle M is a complex manifold equipped with a free holomorphic action of a compact complex torus T . Such a manifold is fibered over M/T , with fiber T . We discuss the notion of positivity in fiber bundles and define positive toric bundles. Given an irreducible complex subvariety $X \subset M$ of a positive principal toric bundle, we show that either X is T -invariant, or it lies in an orbit of T -action. For principal elliptic bundles, this theorem is known ([V]). As follows from Borel-Remmert-Tits theorem, any simply connected compact homogeneous complex manifold is a principal toric bundle. We show that compact Lie groups with left-invariant complex structure I are positive toric bundles, if I is generic. Other examples of positive toric bundles are discussed.

Contents

1	Positivity in fiber bundles	3
1.1	Positive elliptic fibrations	3
1.2	Convexity and positivity	5
2	Homogeneous complex manifolds	8
2.1	Homogeneous complex manifolds and toric fibrations	8
2.2	A generalization of Borel-Remmert-Tits theorem	9
2.3	The Calabi-Eckmann construction	11
2.4	Samelson spaces	12
3	Fiber bundles associated with principal fibrations	14
3.1	Varieties of Fujiki class C	14
3.2	Multisections of fiber bundles and positivity	17
4	Relative Douady spaces	19

¹Misha Verbitsky is an EPSRC advanced fellow supported by EPSRC grant GR/R77773/01

Introduction

Principal toric bundles were always a source of examples in complex geometry (classical Hopf manifolds and Calabi-Eckmann spaces come to mind), but the recent progress in string theory in interaction with algebraic geometry puts a new light on this classical subject. Since a discovery of the hypercomplex structures on compact Lie groups and other homogeneous spaces by the physicists Spindel et al ([SSTV]) and then Joyce ([J]), the complex geometry of homogeneous spaces became a focus of much research, being a target space of supersymmetric string theories with rich supersymmetry. However, the algebraic geometry of these spaces is not much understood. This paper is an attempt to fill this gap, bringing a number of results about subvarieties of principal toric bundles, in particular - compact homogeneous complex manifolds with abelian fundamental group.

A principal toric bundle M is a complex manifold equipped with a free holomorphic action of a compact complex torus T . Such a manifold is fibered over M/T , with fiber T . In the sequel, we discuss positivity in fiber bundles and define positive toric bundles, generalizing a notion developed for elliptic bundles in [V].

Many examples of principal toric bundles appear from a theory of complex homogeneous manifolds. It is known since 1960-es that a simply connected compact complex homogeneous manifold is a principal toric fibration over a rational projective homogeneous manifold (this is known as **the Tits fibration**). In fact, this result holds for a complex homogeneous manifold with abelian fundamental group (Theorem 2.3).

A principal toric fibration obtained this way from a general complex structure on a Lie group is always positive (Theorem 2.7).

We study complex subvarieties of positive principal toric fibrations. Given an irreducible complex subvariety $X \subset M$ of a positive principal toric bundle, we show that either X is T -invariant, or it lies in an orbit of T -action (Theorem 1.13).

This theorem is elementary for $\dim_{\mathbb{C}} T = 1$, but for $\dim_{\mathbb{C}} T > 1$ the proof uses highly non-trivial results on deformation theory and the spaces of Fujiki class C, which are due to Fujiki and Demailly-Paun.

1 Positivity in fiber bundles

1.1 Positive elliptic fibrations

For holomorphic vector bundles and principal elliptic bundles, the subject of positivity is well established. These notions are related geometrically, as we shall explain now.

Let X be a projective manifold, and L a holomorphic line bundle on X . Positivity of L manifests itself in the geometric properties of the total space $\text{Tot}(L^*)$ of the non-zero vectors in the dual bundle, which is a principal $\mathbb{C}^*$ -fibration over X . When L is positive, the projection $\text{Tot}(L^*) \xrightarrow{\pi} X$ has no multisections. Indeed, any such multisection, that is, a subvariety $X' \subset \text{Tot}(L^*)$ which is surjective and finite over X , gives a holomorphic section $\gamma_{X'}$ of $(L^*)^{\otimes N}$, where N is cardinality of $\pi^{-1}(x) \cap X'$ for general $x \in X$. The section $\gamma_{X'}$ is defined as a functional on $(L)^{\otimes N}$, mapping

$$v_1 \otimes v_2 \otimes \dots \otimes v_N \in L^{\otimes N} \Big|_x$$

to

$$\prod_{z_i \in \pi^{-1}(x) \cap X'} v_i(z_i).$$

When L is positive, $(L^*)^N$ has no sections, hence $\text{Tot}(L^*) \xrightarrow{\pi} X$ has no multisections.

For example, when X is $\mathbb{C}P^n$ and L is $\mathcal{O}(1)$, $\text{Tot}(L^*)$ is identified with $\mathbb{C}^{n+1} \setminus 0$. This space is quasi-affine and cannot contain compact subvarieties.

This effect is even more striking if we take a quotient $M := \text{Tot}(L^*) / \langle q \rangle$ of $\text{Tot}(L^*)$ by the $\mathbb{Z}$ -action mapping v to qv , where $q \in \mathbb{C}^*$ is a fixed number, $|q| > 1$. This quotient is a principal elliptic bundle over X , with the fiber isomorphic to $T = \mathbb{C}^* / \langle q \rangle$. When L is positive, this manifold is called a **regular Vaisman manifold** (see e.g. [DO], [OV2]).

It was shown (see e.g. [V]) positive-dimensional complex subvarieties of this principal elliptic fibration are T -invariant whenever L is ample. In other words, for any closed subvariety $Z \subset M$, Z is a preimage of $Z_1 \subset X$, for some complex subvariety of X .

If, in addition, $\dim_{\mathbb{C}} X > 1$, a similar result is true for all stable holomorphic bundles on M , and even all stable reflexive coherent sheaves on M . Namely, given a stable reflexive coherent sheaf F , there exists a line bundle $\mathcal{L}$ on M such that $F \cong \pi^* F_1 \otimes \mathcal{L}$, where F_1 is a stable coherent sheaf on X ([V]).

This result has a number of far-reaching consequences, even in the most trivial example when $X = \mathbb{C}P^n$, $L = \mathcal{O}(1)$ and $M = \mathbb{C}^{n+1} \setminus 0 / \langle q \rangle$ is a classical Hopf manifold. However, the argument of [V] is much more general, and can be applied to other principal elliptic bundles.

Definition 1.1: A **principal elliptic fibration**, or a **principal elliptic bundle** is a complex manifold M equipped with a free holomorphic action of a 1-dimensional compact complex torus T . We consider a principal elliptic fibration as a T -fibration $M \xrightarrow{\pi} M/T$.

Examples of these include the Hopf manifold and regular Vaisman manifolds defined above, but also such interesting objects as a Calabi-Eckmann manifold, that is, a product of two odd-dimensional spheres $S^{2n-1} \times S^{2m-1}$, $m, n > 1$, equipped with an $SO(2n) \times SO(2m)$ -invariant complex structure (see Subsection 2.3). In [V], the positivity of principal elliptic bundles was defined as follows.

Definition 1.2: Let $M \xrightarrow{\pi} X$ be a principal elliptic fibration, with X a Kähler manifold. We call π a **positive elliptic fibration**, if for some Kähler form ω on X , $\pi^*\omega$ is exact.

Of course, all principal elliptic fibrations $M \xrightarrow{\pi} X$ with $H^2(M) = 0$ and X Kähler are necessarily positive. This includes all Hopf manifolds (which are diffeomorphic to $S^1 \times S^{2n-1}$) and the Calabi-Eckmann manifolds $S^{2n-1} \times S^{2m-1}$.

For a manifold $M := \text{Tot}(L^*) / \langle q \rangle$ considered above, positivity is equivalent to the positivity of the corresponding line bundle L .

The positivity of elliptic fibrations has powerful geometric applications.

Theorem 1.3: ([V]) Let X be a compact Kähler manifold, and $M \xrightarrow{\pi} X$ a positive principal elliptic fibration. Then, for any positive-dimensional, irreducible subvariety $Z \subset M$, $Z = \pi^{-1}(Z_1)$, for some complex subvariety $Z_1 \subset X$. If, in addition, $\dim_{\mathbb{C}} X > 1$, then for any stable reflexive coherent sheaf F on M , $F = \pi^*F_1 \otimes \mathcal{L}$, where F_1 is a stable coherent sheaf on X and L a line bundle.

■

1.2 Convexity and positivity

For the purposes of the present paper, a good notion of positivity in fiber bundles would give the same consequences as in Theorem 1.3. All irreducible subvarieties of a positive fiber bundle $M \xrightarrow{\pi} X$ should lie in the fiber or have the form $\pi^{-1}(Z)$; and all stable bundles must be pullbacks, up to tensoring with a line bundle.

Further on in this paper, we study principal toric fibrations, that is, fibrations with a fiber which is a compact complex torus. They are defined in the same way as principal elliptic fibrations.

Definition 1.4: Let T be a compact complex torus. A **principal toric fibration**, or a **principal toric bundle** with a fiber T is a complex manifold M equipped with a free holomorphic action of T . We consider a principal toric fibration as a T -fibration $M \xrightarrow{\pi} M/T$.

Principal toric fibrations appear in different disguises in all kinds of algebro-geometric research. A classification of these objects is found in [Hö].

Even for principal toric fibrations, it is not clear what is a good notion of positivity. One could require that $\pi^*\omega$ is exact, for some Kähler form ω on X . This is clearly not enough. Indeed, one could take a positive elliptic fibration, and its product with a torus T_1 , as a toric fibration. The pullback $\pi^*\omega$ is exact. However, any subvariety of $X \times T_1$ (and there could be many) gives a corresponding subvariety of M .

Still, this notion has useful geometric consequences.

Definition 1.5: Let $M \xrightarrow{\pi} X$ be a fibration of complex manifolds. We say that π is **convex** if for some Kähler form ω on X , $\pi^*\omega$ is exact.

Definition 1.6: Let $M \xrightarrow{\pi} X$ be a fibration of complex manifolds, and $X' \subset M$ a closed subvariety. Assume that the restriction $\pi|_{X'} : X' \rightarrow X$ is proper, surjective and generically finite. Then X' is called a **multisection** of π .

The following trivial lemma explains the utility of convexity in fiber bundles.

Lemma 1.7: Let $M \xrightarrow{\pi} X$ be a morphism of complex manifolds, X compact and Kähler, ω a Kähler class. Assume that $\pi^*\omega$ is exact. Then π

does not admit multisections.

Proof: Let X' be a multisection, with a generic point $x \in X$ having precisely N pre-images in X' under the map $\pi|_{X'} : X' \rightarrow X$. Then

$$\int_{X'} \pi^* \omega^{\dim X} = N \int_X \omega^{\dim X} > 0.$$

This is impossible, because $\pi^* \omega$ is exact. ■

The same argument is valid in weaker assumptions.

Definition 1.8: Let M be a complex manifold, $\dim_{\mathbb{C}} M = n$, and ω a $(1,1)$ -form, which is positive and closed. We say that ω is **weakly Kähler** if $\omega^n \neq 0$ somewhere on M . Of course, in this case $\int_M \omega^n > 0$ (assuming ω^n has compact support).

Definition 1.9: Let $M \xrightarrow{\pi} X$ be a fibration of complex manifolds. We say that π is **weakly convex** if for some weakly Kähler form ω on X , $\pi^* \omega$ is exact.

Remark 1.10: Clearly, Lemma 1.7 is valid in assumption of weak convexity.

Let $M \xrightarrow{\pi} X$ be a principal toric bundle with fiber T , and V a complex variety equipped with a holomorphic action of T . Consider the quotient $M_V := (M \times V)/T$, with T acting diagonally on M and V . Clearly, $M_V \xrightarrow{\pi_V} X$ is a locally trivial holomorphic fibration over X , with fiber V .

Definition 1.11: In this situation, $M_V \rightarrow X$ is called **a fiber bundle associated with a principal toric bundle $M \rightarrow X$** .

For principal toric bundles, the following notion of positivity seems to be useful.

Definition 1.12: Let $M \xrightarrow{\pi} X$ be a principal toric bundle with fiber T . Assume that for all surjective homomorphisms $T \rightarrow T_1$ from T to a positive-dimensional torus T_1 , the associated bundle M_{T_1} is (weakly) convex. Then $M \xrightarrow{\pi} X$ is called (weakly) **Q-positive**.

The main result of this paper is the following theorem

Theorem 1.13: Let $M \xrightarrow{\pi} X$ be a principal toric bundle, and $Z \subset M$ an irreducible complex subvariety. Assume that π is compact and Q-positive. Then Z is T -invariant¹, or Z lies in a fiber of π .

Proof: We prove Theorem 1.13 in Section 4. ■

Remark 1.14: For $\dim T = 1$, this result follows directly from Lemma 1.7.

Remark 1.15: For $\dim T > 1$, we do not have any control over the category of coherent sheaves on a positive T -fibration. It is not clear whether Q-positivity will imply results similar to Theorem 1.3, showing that all stable bundles are pullbacks (up to a multiplication by a line bundle). This is the reason for “Q-” in Q-positivity (“Q” stands for “quotient”).

Remark 1.16: For examples of Q-positive toric fibrations, see Theorem 2.7.

Remark 1.17: Let $M \xrightarrow{\pi} X$ be a principal toric bundle, and $X_1 \rightarrow X$ a holomorphic bimeromorphic map. Then $M \times_X X_1 \xrightarrow{\pi_1} X_1$ is a principal toric bundle. When π is Q-positive, π_1 is not necessarily Q-positive; however, it is weakly Q-positive. Similarly, if $Z \subset X$ is a closed subvariety, $Z_1 \rightarrow Z$ its resolution of singularities, and $M_1 := \pi^{-1}(Z) \times_Z Z_1 \xrightarrow{\pi_1} Z_1$ the corresponding principal toric bundle, π_1 is weakly Q-positive, though not necessarily Q-positive.

Recall that a compact complex torus T is called **simple** if for any holomorphic homomorphism $T \xrightarrow{\varphi} T_1$ to another torus, $\ker \varphi$ is finite. A generic complex torus is simple; a generic abelian variety is also simple. Actually, non-simple tori are quite rare, and the codimension of the space of non-simple tori inside the moduli space of all tori grows linearly with the dimension of the torus; same is true for abelian manifolds.

Claim 1.18: Let $M \xrightarrow{\pi} X$ be a principal toric bundle with fiber T . Assume that T is simple and M is (weakly) convex. Then M is (weakly) Q-positive.

Proof: Let ω be a (weakly) Kähler form on X such that $\pi^*\omega = d\theta$ is exact, and $T \xrightarrow{\mu} T_1$ a surjective homomorphism to a positive-dimensional

¹That is, takes a form $\pi^{-1}(Z_1)$ for some $Z_1 \subset X$

torus T_1 . Since T is simple, μ is an isogeny (a finite quotient map). Therefore, $M_{T_1} \xrightarrow{\pi_1} X$ is a quotient of M over a finite group $\Gamma = \ker \mu$ freely acting on M . Averaging θ with Γ , we obtain a 1-form θ_Γ on M_{T_1} which satisfies $d\theta_\Gamma = \pi_1^* \omega$. ■

2 Homogeneous complex manifolds

2.1 Homogeneous complex manifolds and toric fibrations

This paper appeared as an attempt to understand the algebraic geometry of compact, homogeneous, simply connected manifolds. These manifolds were classified by H. C. Wang ([Wa]) and much studied during 1950-ies and the 1960-ies. H. Samelson ([Sa]) discovered homogeneous complex structures on even-dimensional semisimple Lie groups. In 1980-es and 1990-ies the Samelson manifolds became more and more important in string physics, differential geometry and algebraic geometry.

Ever since the physicists Spindel et al ([SSTV]) and then Joyce ([J]) constructed homogeneous hypercomplex structures on some compact Lie groups, generalizing Samelson's work, these manifolds became a favourite testbed for all kinds of conjectures and constructions in quaternionic geometry and sometimes in physics (see e.g. [OP]). In addition, these manifolds are examples of two rather unique (and mysteriously related) geometries, which arose in string theory. The Joyce's hypercomplex manifolds are the only known in dimension > 4 examples of so-called *strong HKT geometry* ([HP], [GP]). In addition, these manifolds are among the few compact examples of *generalized Kähler manifolds* (known in physics as *manifolds with (2,2)-supersymmetry*); see [GHR], [Hi]. Combining these two structures, one obtains an even more mysterious (and rare) geometry, called *generalized hyperkähler geometry*, or (4,4)-supersymmetry ([GHR], [Br], [MV]), which also occurs on Joyce's hypercomplex manifolds.

From their construction, which is explained in Subsection 2.3, it is clear that the Samelson manifolds are fibered over a flag space, with fibers being compact tori. It is a special case of a general result due to Borel and Remmert ([BR]) and Tits ([T]), and its generalization, proven in this paper (Theorem 2.3); see also [GR].

Theorem 2.1: Let M be a compact, homogeneous, simply connected complex manifold. Then M is a principal toric fibration over a base X , which is a homogeneous, rational, projective manifold.

Proof: Recall that a complex manifold is called **parallelizable** if its holomorphic tangent bundle is trivial. The following lemma is used.

Lemma 2.2: ([T]) Let M , $\dim_{\mathbb{C}} M = n$ be a compact homogeneous complex manifold with trivial canonical bundle. Then M is parallelizable.

Proof: The holomorphic tangent bundle TM is globally generated. Chose a set $\gamma_1, \dots, \gamma_n$ of holomorphic vector fields which are linearly independent at $m \in M$. Since the canonical bundle of M is trivial, $\gamma_1, \dots, \gamma_n$ are linearly independent everywhere. ■

Return to the proof of Theorem 2.1. Since TM is globally generated, the anticanonical bundle $K^{-1}M$ is globally generated too, hence base point free. This gives a holomorphic morphism $M \xrightarrow{\psi} \mathbb{P}(\Gamma(K^{-1}M))$ from M to the projectivization of the space of sections of $K^{-1}M$. This map is obviously equivariant with respect to the action of the group $G = \text{Aut}(M)$. Therefore, the projective manifold $B = \psi(M)$ is homogeneous, and hence rational. By adjunction formula, the fibers F of ψ have trivial canonical class; Lemma 2.2 implies that they are parallelizable. Using the homotopy exact sequence for a fibration

$$\pi_2(B) \longrightarrow \pi_1(F) \longrightarrow \pi_1(M) = 0,$$

we obtain that the fundamental group of F is abelian. Therefore, F is a complex torus (see the proof of Theorem 2.3 below for a detailed argument). The sheaf $T_{\psi}M$ of holomorphic vector fields tangent to the fibers of ψ is a trivial bundle, by the same reasoning as used to prove Lemma 2.2. Let $G_0 \subset \text{Aut}(M)$ be a Lie group generated by the sections of $T_{\psi}M$. Identifying G with the group of automorphisms of fibers of ψ , we obtain that G is a compact torus, and $\psi : M \longrightarrow X$ is a principal toric fibration. ■

2.2 A generalization of Borel-Remmert-Tits theorem

Borel-Remmert-Tits theorem has a generalization, which is quite useful in our work.

Theorem 2.3: Let M be a compact complex manifold equipped with a transitive holomorphic action of a Lie group G . Assume that M admits a G -invariant decomposition $M = M_0 \times (S^1)^n$, where $\pi_1(M_0) = 0$. Let

$\psi : M \longrightarrow B \subset \mathbb{P}(H^0(K^{-1}))$ be the G -equivariant fibration constructed in the proof of Theorem 2.1. Then ψ is a principal toric fibration.

Proof: As we have shown, the fibers F of π are compact, homogeneous manifolds which are holomorphically parallelizable, that is, have trivial tangent bundle. To prove Theorem 2.3 we have to show that these fibers are complex tori. It is well known (see [Wo]) that such manifolds are always of form $F = \mathcal{G}/\Gamma$, where $\mathcal{G}$ is a complex Lie group, and Γ a cocompact lattice. To prove that F is a torus, it suffices to show that $\pi_1(F)$ is abelian. Indeed, Γ is a subgroup of $\pi_1(M)$, and (being a cocompact lattice), Γ is Zariski dense in $\mathcal{G}$, hence $\mathcal{G}$ is also abelian. We reduced Theorem 2.3 to the following lemma.

Lemma 2.4: In assumptions of Theorem 2.3, we have a decomposition $F = F_0 \times (S^1)^n$, where $\pi_1(F_0) = \pi_2(B)/j(\pi_2(M))$, where $j : \pi_2(M) \longrightarrow \pi_2(B)$ is the map induced by the projection.

Proof: Since G preserves the decomposition $M = M_0 \times (S^1)^n$, G contains a torus $T^n = (S^1)^n$ acting trivially on M_0 and tautologically on $(S^1)^n$. Consider the natural action of $T^n \subset G$ on B . If any of its orbits has positive dimension, it is isomorphic to a quotient $T = T^n/T_0$. This gives a free holomorphic action of a torus T on B . This is impossible, because B is rational, and any holomorphic vector field on B vanishes somewhere. We have shown that T^n acts trivially on B , and the map $\psi : M \longrightarrow B$ factors through the standard projection $M = M_0 \times (S^1)^n \xrightarrow{\pi} M_0$. Therefore, $F = F_0 \times T^n$, with F_0 being a fiber of the standard fibration $M_0 \longrightarrow B$. To see that the fundamental group of F_0 is abelian, we use an exact sequence of homotopy

$$\pi_2(M_0) \xrightarrow{j} \pi_2(B) \longrightarrow \pi_1(F_0) \longrightarrow \pi_1(M_0) = 0.$$

associated to the fibration $M_0 \longrightarrow B$. This gives an isomorphism $\pi_1(F_0) = \pi_2(B)/j(\pi_2(M))$. Lemma 2.4 is proven. We proved Theorem 2.3. ■

Remark 2.5: This theorem has a generalization, also due to J. Tits ([T]). Let M be a compact homogeneous manifold, with a Lie group G acting on M transitively. Then M is fibered over a projective, rational, homogeneous space with a fiber which is homogeneous and parallelizable. This fibration is called **the Tits fibration**. Tits also proved that it is G -invariant, and universal.

Quite often, a compact, homogeneous complex manifold would have $H^2(M, \mathbb{Q}) = 0$. The Calabi-Eckmann manifolds and the Hopf manifolds we mentioned previously clearly satisfy $H^2(M) = 0$. Let G be a compact Lie group with $H^1(G, \mathbb{Q}) = 0$. The cohomology algebra of G is a Hopf algebra; by Hopf theorem ([Ho]), $H^*(G, \mathbb{Q})$ is a free exterior algebra, with generators of odd degree. In particular, $H^2(G, \mathbb{Q}) = 0$.

If G is even-dimensional, G admits a left-invariant complex structure; if G is odd-dimensional, $G \times S^1$ admits a left-invariant complex structure. These manifolds are called *Samelson spaces* (see Subsection 2.4). In both of these cases, $H^2(M, \mathbb{Q}) = 0$, and the corresponding principal toric bundle (Theorem 2.3) is convex. If the torus T is, in addition, simple, M is $\mathbb{Q}$ -positive. Further on in this section, we explain that this happens for generic complex structures on G and $G \times S^1$.

2.3 The Calabi-Eckmann construction

The complex structures on the Lie groups can be constructed explicitly, in such a way that the principal toric action becomes apparent. To see this we first review the construction of Calabi-Eckmann manifolds.

Consider the principal $\mathbb{C}^* \times \mathbb{C}^*$ -fibration

$$\tilde{M} = \mathbb{C}^n \setminus 0 \times \mathbb{C}^m \setminus 0 \longrightarrow X = \mathbb{C}P^{n-1} \times \mathbb{C}P^{m-1}.$$

Given a subgroup $G \subset \mathbb{C}^* \times \mathbb{C}^*$, we obtain an associated fiber bundle $M_G = \tilde{M}/G \longrightarrow X$, with a fiber $\mathbb{C}^* \times \mathbb{C}^*/G$. We parametrize $\mathbb{C}^* \times \mathbb{C}^*$ by the set of pairs $e^{t_1} \times e^{t_2}$, $t_1, t_2 \in \mathbb{C}$. Given a number $\alpha \in \mathbb{C} \setminus \mathbb{R}$, we observe that the subgroup

$$G := \{e^t \times e^{\alpha t} \in \mathbb{C}^* \times \mathbb{C}^*, \quad t \in \mathbb{C}\} \subset \mathbb{C}^* \times \mathbb{C}^*$$

is co-compact, and the quotient $\mathbb{C}^* \times \mathbb{C}^*/G$ is an elliptic curve T , isomorphic to the quotient

$$\mathbb{C}/\langle 2\pi\sqrt{-1}n + 2\pi\alpha\sqrt{-1}m \rangle, \quad n, m \in \mathbb{Z}$$

Therefore, the fibration $M_G \longrightarrow X$ is an elliptic fibration, with fiber T ; it is easy to check that M_G is actually diffeomorphic to $S^{2n-1} \times S^{2m-1}$. We have constructed the Calabi-Eckmann manifold.

A similar construction will work whenever we have a principal $(\mathbb{C}^*)^{2n}$ -bundle $\tilde{M} \longrightarrow X$. We may identify $(\mathbb{C}^*)^{2n}$ with a quotient of $W = \mathbb{C}^{2n}$ by a subgroup

$$\Gamma = \left\{ \sum_{i=1}^{2n} 2\pi\sqrt{-1} k_i \mid k_i \in \mathbb{Z} \right\}$$

isomorphic to $\mathbb{Z}^{2n}$. Choosing an n -dimensional complex subspace $V \subset W$ not intersecting $\mathbb{R}\Gamma$, we find that the quotient $T = W/\langle V + \Gamma \rangle$ is a compact torus (this is actually a standard construction of compact tori, used in their classification).

Now consider V as a subgroup of $(\mathbb{C}^*)^{2n} = \mathbb{C}^{2n}/\Gamma$. Clearly, the quotient $\tilde{M}/V$ is a principal toric bundle over X , with fiber T .

2.4 Samelson spaces

The left-invariant complex structures on Lie groups were constructed by H. Samelson in 1954 ([Sa]).

Definition 2.6: Let G be an even-dimensional, compact, semisimple Lie group, and I a left-invariant complex structure on G . Then (G, I) is called a **Samelson space**. It is a homogeneous complex manifold.

Let G be an even-dimensional compact Lie group; it is obviously semisimple. Denote by $G_{\mathbb{C}}$ its complexification, and let $B \subset G_{\mathbb{C}}$ be its Borel subgroup, and $F = G_{\mathbb{C}}/B$ the corresponding flag space, which is obviously compact. The group G can be considered as a subgroup in $G_{\mathbb{C}}$; the natural projection $G \rightarrow F$ is surjective, and has the maximal torus of G as its fiber. This is the toric fibration we require; it remains only to produce a complex structure on G which is compatible with this fibration.

Denote by U the unipotent radical of B . The space $\tilde{M} := G_{\mathbb{C}}/U$ is fibered over F with a fiber $T_{\mathbb{C}} = (\mathbb{C}^*)^k$, with $k = \dim_{\mathbb{R}} G - 2\dim_{\mathbb{C}} F$ being even. We arrive at the situation described at the end of Subsection 2.3: a $(\mathbb{C}^*)^{2n}$ -fibration over a compact base. Taking a quotient of $\tilde{M}$ with a subgroup $V \subset T_{\mathbb{C}} =$ isomorphic to $\mathbb{C}^n$, as indicated in Subsection 2.3, we obtain a principal toric fibration $M = \tilde{M}/V$. Embedding G into $\tilde{M} = G_{\mathbb{C}}/U$, and projecting to M , we identify G with M , and this identification is by construction G -invariant.

Clearly, this principal fibration is the one postulated by Borel-Remmert-Tits theorem (Theorem 2.1).

Notice that we have a complete freedom of choosing $V \subset T_{\mathbb{C}}$. Therefore the left-invariant complex structure on G can be chosen in such a way that any given torus of appropriate dimension becomes a fiber of the Tits fibration.

There is another construction of Samelson manifolds, which is even more elementary and explicit. Take the Lie algebra $\mathfrak{g}$ of a compact, even-dimensional Lie group G , and let $\mathfrak{g}_{\mathbb{C}}$ be its complexification. Take a positive

root system $e_1, \dots, e_n \in \mathfrak{g}_{\mathbb{C}}$, and let $f_1, \dots, f_n$ the corresponding negative roots, and $\mathfrak{h}_{\mathbb{C}} \subset \mathfrak{g}_{\mathbb{C}}$ be the corresponding Cartan subalgebra, so that

$$\mathfrak{g}_{\mathbb{C}} = \langle e_1, \dots, e_n, f_1, \dots, f_n \rangle \oplus \mathfrak{h}_{\mathbb{C}}.$$

Let $\mathfrak{h}$ be a Lie algebra of a maximal torus of G such that $\mathfrak{h}_{\mathbb{C}}$ is its complexification. The space $\mathfrak{g}$ is decomposed as

$$\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{F},$$

where $\mathfrak{F}$ is the space of real vectors in $\langle e_1, \dots, e_n, f_1, \dots, f_n \rangle$. Since $\dim G$ is even, the real vector space $\mathfrak{h}$ is even-dimensional.

Now choose a complex structure $\mathcal{I}$ on $\mathfrak{g}$, in such a way that the decomposition $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{F}$ is preserved. To define a complex structure $\mathcal{I}$ on $\mathfrak{F}$ it is the same as to define a decomposition

$$\mathfrak{F} \otimes \mathbb{C} = \mathfrak{F}^{1,0} \oplus \mathfrak{F}^{0,1},$$

with $\overline{\mathfrak{F}^{1,0}} = \mathfrak{F}^{0,1}$, and $\mathcal{I}$ acting as $\sqrt{-1}$ on $\mathfrak{F}^{1,0}$ and as $-\sqrt{-1}$ on $\mathfrak{F}^{0,1}$. We define

$$\mathfrak{F}^{1,0} = \langle e_1, \dots, e_n \rangle, \quad \mathfrak{F}^{0,1} = \langle f_1, \dots, f_n \rangle.$$

Choose the action of $\mathcal{I}$ on $\mathfrak{h}$ arbitrarily.

Using the left action of G on itself, $\mathcal{I}$ becomes a left-invariant almost complex structure I . By Newlander-Nirenberg, I is integrable if and only if $[T^{0,1}G, T^{0,1}G] \subset T^{0,1}G$. This is equivalent to

$$[\mathfrak{g}^{0,1}, \mathfrak{g}^{0,1}] \subset \mathfrak{g}^{0,1},$$

where $\mathfrak{g}^{0,1} \subset \mathfrak{g}_{\mathbb{C}}$ is a space of all vectors where $\mathcal{I}$ acts as $-\sqrt{-1}$. However, from the construction of $\mathcal{I}$ it is apparent that $\mathfrak{g}^{0,1}$ is identified with a subspace of Cartan algebra plus $\langle f_1, \dots, f_n \rangle$, and such a space is closed under commutators.

It is not difficult to check that this construction is in fact identical to that we gave earlier. The same argument was used by Samelson to construct an invariant complex structure on G/T , where G is a compact Lie group and $T \subset G$ a compact torus of even codimension in G .

We conclude this section with the following theorem, which follows from the constructions we have given.

Theorem 2.7: Let G be a compact, even-dimensional Lie group, and I a left-invariant complex structure on G . Consider the corresponding Tits

fibration $(G, I) \xrightarrow{\pi} F$, with fibers identified with a compact complex torus T . Assume that T is simple¹. Then the fibration π is $\mathbb{Q}$ -positive.

Proof: Let

$$G = G_0 \times S_1^k,$$

where G_0 is a semisimple group with $H^1(G_0, \mathbb{Q}) = 0$. Such a decomposition follows from the classification of compact Lie groups. It is easy to see that G is a toric fibration over the flag space F associated with G_0 ([Sa]). However, $H^2(G_0, \mathbb{Q}) = 0$ as we have mentioned above. The Tits map $G \xrightarrow{\pi} F$ is a composition of the projection $G \rightarrow G_0$ and the natural map $G_0 \rightarrow F$. Therefore, the pull-back homomorphism

$$\pi^* : H^2(F, \mathbb{R}) \rightarrow H^2(G, \mathbb{R})$$

vanishes. We obtain that $\pi^*\omega$ is exact for any Kähler form ω on F . This implies that (G, I) is convex. Using Claim 1.18, we obtain that (G, I) is $\mathbb{Q}$ -positive. ■

Remark 2.8: The same argument works to show that for any torus $T \subset G$ of even codimension in a compact Lie group G , a generic left-invariant complex structure on G/T is $\mathbb{Q}$ -positive. Such spaces were constructed and classified by Samelson in [Sa].

3 Fiber bundles associated with principal fibrations

3.1 Varieties of Fujiki class C

Definition 3.1: Let X, Y be complex varieties, and $C \subset X \times Y$ a closed subvariety, such that the projections $C \rightarrow X, C \rightarrow Y$ are proper and generically one-to-one. Such C is called a **bimeromorphic morphism**, or **bimeromorphic modification** between X and Y .

Clearly, a composition of bimeromorphic maps is well defined, and is again bimeromorphic.

Definition 3.2: Let M be a complex variety. We say that M is in **Fujiki class C** if M is bimeromorphic to a Kähler manifold.

¹This happens if I is generic, as shown above.

Varieties of Fujiki class C were defined by A. Fujiki in [F1], and explored at great length in subsequent papers ([F2], [F3]).

The following properties of Fujiki class C will be useful for our work.

Proposition 3.3: (Fujiki) Let M be a variety of Fujiki class C, and $X \subset M$ a closed complex subvariety. Then X is also of Fujiki class C.

Proof: [F2], Lemma 4.6 ■

Theorem 3.4: Let M be a compact variety of Fujiki class C, $X \subset M$ a closed subvariety, and D its Douady deformation space. Then D is compact and belongs to Fujiki class C.

Proof: [F2], [F3]. ■

In [DP], J.-P. Demailly and M. Paun gave a simple characterization of Fujiki class C.

Recall that a **p -current** on a manifold M is a p -form taking values in distributions. We consider currents as functionals on smooth $(\dim_{\mathbb{R}} M - p)$ -forms with compact supports, called **test-forms**. De Rham differential on p -currents is defined by the formula

$$\langle d\eta, \psi \rangle := (-1)^p \langle \eta, d\psi \rangle,$$

where ψ is a test-form, and η is a p -current. The cohomology of currents are equal to de Rham cohomology of the manifold. The Hodge decomposition on currents

$$\mathcal{C}^p(M) = \bigoplus_{i+j=p} \mathcal{C}^{i,j}(M)$$

is defined in the same manner. A $(1,1)$ -current η is called **positive** if $\langle \eta, \psi \rangle \geq 0$ for any test-form $\psi \in \Lambda^{\dim_{\mathbb{R}} M - 1, \dim_{\mathbb{R}} M - 1}(M)$ which is positive, that is, Hodge dual to a pseudo-Hermitian $(1,1)$ -form with non-negative eigenvalues.

A **closed current** is a current η satisfying $d\eta = 0$. A **Kähler current** is a closed $(1,1)$ -current η such that there exists a Hermitian form ω on M with $\eta - \omega$ is positive.

The cohomology classes of non-smooth Kähler currents can be non-Kähler. For instance, take a divisor $Z \subset M$. The divisor Z can be considered as a $(1,1)$ -current by taking

$$\langle Z, \psi \rangle := \int_Z \psi.$$

If Z is an exceptional divisor, and ω a Kähler form, $\lambda \in \mathbb{R}^{\geq 0}$, the current $\omega + \lambda Z$ is Kähler, but for λ sufficiently big, its cohomology class is clearly non-Kähler.

Demailly and Paun proved the following theorem.

Theorem 3.5: ([DP, Theorem 3.4]) Let M be a compact complex manifold. Then M lies in Fujiki class C if and only if M supports a Kähler current.

■

Remark 3.6: Such a Kähler current is easy to construct. Given a bimeromorphism $M \sim M_1$, where M_1 is Kähler, and using Hironaka desingularization theorem, we can construct a resolution M_2 which dominates M and M_1 and is obtained by sequence of blow-ups with smooth centers. A blow-up of a Kähler manifold is again Kähler, hence M_2 is Kähler. Consider the map $M_2 \xrightarrow{\pi} M$; it is bimeromorphic and holomorphic. Let ω be a Kähler form on M_2 . We consider ω as a current $\underline{\omega}$ on M using the formula

$$\langle \underline{\omega}, \psi \rangle := \langle \omega, \pi^* \psi \rangle,$$

This current is obviously positive and Kähler.

Further on, we shall need the following proposition.

Proposition 3.7: Let M be a compact complex variety of Fujiki class C and G a compact Lie group acting on M holomorphically. Then M admits a G -equivariant, smooth Kähler resolution of singularities, with G -invariant Kähler metric.

Proof: Without restricting the generality, we may assume that M is smooth. In this case, property C is equivalent to existence of a Kähler current. Averaging with G , we may assume that this Kähler current is G -invariant. The proof of Theorem 3.5 involves approximating a given Kähler current by Kähler currents with controlled logarithmic singularities at Z_k , and blowing up the corresponding singular sets ([DP, Theorem 3.2]). The sets Z_k are obtained explicitly as zeroes of multiplier ideal sheaves with prescribed weights on M ([D]). These weights can be chosen G -invariant, because (as seen from the proof) the corresponding weights ψ_k satisfy $|\psi_k - g\psi_k| < C$, for any $g \in G$. This implies that Z_k can be chosen G -invariant. Therefore, the Kähler blow-up of M can be achieved by blowing up the G -invariant subvarieties. This implies existence of a G -equivariant Kähler

resolution of M . Averaging the Kähler form with G , we obtain a G -invariant Kähler metric. ■

3.2 Multisections of fiber bundles and positivity

Recall that a **multisection** of a fiber bundle $M_V \xrightarrow{\pi_V} X$ is an irreducible subvariety $X' \hookrightarrow M_V$ which is proper and finite over a general point of X .

The main result of this section is the following theorem, which ensures that under appropriate conditions a fiber bundle associated with a principal toric bundle does not admit multisections.

Theorem 3.8: Let $M \xrightarrow{\pi} X$ be a principal toric bundle, with fiber T , and V a compact complex variety of Fujiki class C, equipped with a holomorphic T -action. Consider the associated fiber bundle $M_V \xrightarrow{\pi_V} X$, $M_V = M \times V/T$, and let $X' \hookrightarrow M_V$ be its multisection. Assume that M is compact and weakly Q -positive. Then X' is T -invariant.

Proof:

Step 1. Without restricting the generality, we may assume that X' intersects the smooth part of M_V . Indeed, if X' lies in the singular part V_{sing} of V , we may replace M_V with $M_{V_{sing}}$. If, again, X' lies in the singular part of V_{sing} , we replace V_{sing} with its singular part and so on. Using induction by $\dim V$, we obtain that at some step we arrive at the situation when X' intersects the non-singular part.

Step 2. Let $T_0 \subset T$ be a stabilizer $St(z)$ of a generic point $z \in X'$. If $T = T_0$, Theorem 3.8 is proven. Replacing T by T/T_0 , and V by its subset fixed by T_0 , we arrive at assumptions of Theorem 3.8; however, now $St(z) = 0$ for generic $z \in X'$. It suffices to prove Theorem 3.8 in this assumption.

Step 3. Let V' be a T -equivariant Kähler resolution of singularities of V , which exists by Proposition 3.7. A proper preimage of X' under the natural map $M_{V'} \rightarrow M_V$ is a multisection of $M_{V'} \xrightarrow{\pi_{V'}} X$. Hence it suffices to prove Theorem 3.8 assuming that V is Kähler and smooth.

Step 4. For any holomorphic action of a torus T on a complex manifold Z , denote by $St_0(z) \subset T$ the connected component of the stabilizer of $z \in Z$,

and by T_0 the connected component of the group

$$\{t \in T \mid t \text{ acts trivially on } Z\}.$$

Then $St_0(z) = T_0$. Indeed, $St_0(z)/T_0$ acts faithfully on $Sym^*(T_z Z)$, and this gives a holomorphic embedding of $St_0(z)/T_0$ into an algebraic group, which is affine, and hence Stein. Since $St_0(z)$ is holomorphic and compact, such an embedding is necessarily trivial.

In Steps 1-3, we have reduced Theorem 3.8 to the case when V is Kähler and smooth, and the action of T is free generically. Because of Step 4, it remains now to prove Theorem 3.8 assuming that V is Kähler and smooth, and for any $v \in V$, the stabilizer $St(v)$ is finite.

Now Theorem 3.8 can be deduced from the following lemma.

Lemma 3.9: Let $M \xrightarrow{\pi} X$ be a principal toric bundle with fiber T , and V – a Kähler manifold equipped with an action of T . Assume that X is compact, M is weakly convex, and for any $v \in V$, the stabilizer $St(v)$ is finite. Consider the associated fibration $M_V \xrightarrow{\pi_V} X$. Then π_V admits no multisections.

Proof: Let ω be a weakly Kähler form on X such that

$$\pi^* \omega = d\theta \tag{3.1}$$

is exact. Clearly, to show that π_V admits no multisections, it suffices to prove that $\pi_V^* \omega$ is exact (see Lemma 1.7, Remark 1.10).

Averaging θ over T , we may assume that θ is T -invariant. Denote by $\mathfrak{t}$ the Lie algebra of T . For all $t \in \mathfrak{t}$,

$$d(\theta \lrcorner t) = \text{Lie}_t \theta - (d\theta) \lrcorner t = 0.$$

Therefore, $\langle \theta, t \rangle = \theta \lrcorner t$ is constant. Denote by $\theta^\sim \in \mathfrak{t}^*$ the corresponding form on $\mathfrak{t}$.

Since T is a complex Lie group acting holomorphically, the natural embedding¹ $\mathfrak{t} \xrightarrow{\rho} TV$ is $\mathbb{C}$ -linear. Therefore, the Kähler form ω_V of V is non-degenerate on $\rho(\mathfrak{t})$. Since ω_V is closed and T -invariant, for all $t_1, t_2 \in \mathfrak{t}$, the function $\omega_V(t_1, t_2) \in C^\infty V$ is constant (the argument is the same as used above to show that $\theta \lrcorner t$ is constant). This defines a symplectic form $\omega_V^\sim$ on $\mathfrak{t}$. Let $t_\theta \in \mathfrak{t}$ be defined by the formula

$$\langle \theta^\sim, t \rangle = \omega_V^\sim(t_\theta, t).$$

¹This map is an embedding, because the stabilizer $St(v)$ is finite everywhere. This is the only place we use this assumption in the whole argument.

Denote by θ_V the 1-form $\omega(t_\theta, \cdot)$ on V . By Cartan's formula, θ_V is closed. Moreover,

$$\langle \theta_V, t \rangle = \theta^\vee(t) = \langle \theta, t \rangle \quad (3.2)$$

for all $t \in \mathfrak{t}$.

Consider the projection maps

$$\begin{array}{ccc} & M \times V & \\ \pi_1 \swarrow & & \searrow \pi_2 \\ M & & V \end{array}$$

and let $\Psi := \pi_1^* \theta - \pi_2^* \theta_V$. Since θ_V is closed, (3.1) gives

$$d\Psi = \pi_1^* \pi^* \omega. \quad (3.3)$$

By construction, Ψ is T -invariant; from (3.2), we obtain that $\langle \Psi, t \rangle = 0$ for all $t \in \mathfrak{t}$, with T acting on $M \times V$ diagonally. From T -invariance and $\langle \Psi, t \rangle = 0$ it follows that Ψ is a pullback of a form ψ on $(M \times V)/T = M_V$. Now, (3.3) implies that $\pi_V^* \omega = d\psi$. We proved that $\pi_V^* \omega$ is exact. This finishes the proof of Lemma 3.9 and Theorem 3.8. ■

4 Relative Douady spaces

Using Theorem 3.8, we can prove Theorem 1.13 as follows.

Let $M \xrightarrow{\pi} X$ be a $\mathbb{Q}$ -positive toric fibration, and $Z \subset M$ an irreducible complex subvariety, which does not lie in a fiber of π . Denote by X_1 the resolution of singularities of $\pi(Z)$, and let $M_1 := X_1 \times_{\pi(Z)} \pi^{-1}(\pi(Z)) \xrightarrow{\pi_1} X_1$ be the corresponding principal toric fibration, and Z_1 be the proper preimage of Z in M_1 . Clearly, π_1 is a weakly $\mathbb{Q}$ -positive fibration (see Remark 1.17).

We have reduced Theorem 1.13 to the following statement.

Proposition 4.1: Let $M \xrightarrow{\pi} X$ be a principal toric bundle, $\dim X > 0$, and $Z \subset M$ an irreducible complex subvariety. Assume that π is weakly $\mathbb{Q}$ -positive, and $\pi(Z) = X$. Then $Z = M$.

Proof: Let X be a complex variety, $Y \subset X$ a compact subvariety. In this situation, a space of deformations of Y inside X is defined, called **the Douady space** ([Do], [F2]).¹ Denote by D_Y the component of the Douady

¹There are actually two different definitions of the moduli of deformations of subvarieties, the Douady space and the Barlet space, [Ba]. They are bimeromorphically equivalent. For our purposes any of these two can serve.

space containing Y . In [F3], A. Fujiki proved that if X is a compact manifold of Fujiki class C, same is true for D_Y .

Let D be a Douady space of subvarieties in T , equipped with a natural T -action, and $D_\pi := M_D$ the associated fiber bundle. We call D_π **the relative Douady space of $M \xrightarrow{\pi} X$** . It is a classifying space of subvarieties of the fibers of π .

For $x \in X$, let $Z_x := \pi^{-1}(x) \cap Z$ be the subvariety of a torus $\pi^{-1}(x)$ corresponding to x . This gives a multisection $X' \subset D_\pi$, where D_π is the relative Douady space. Applying Theorem 3.8, we find that any multisection of D_π is T -invariant. This means that $Z_x \subset \pi^{-1}(x)$ is T -invariant. However, the only T -invariant subvariety of a torus T is T . Therefore, $Z = M$. ■

Acknowledgements: I am grateful to J.-P. Demailly for valuable advice and consultations, and to the referee for error-checking and simplification of the proof of Theorem 3.8.

References

- [Ba] Barlet, Daniel *Espace analytique réduit des cycles analytiques complexes compacts d'un espace analytique complexe de dimension finie*, Fonctions de plusieurs variables complexes, II (Sém. François Norguet, 1974–1975), pp. 1–158. Lecture Notes in Math., Vol. 482, Springer, Berlin, 1975.
- [BR] Borel, A.; Remmert, R. *Über kompakte homogene Kählersche Mannigfaltigkeiten*, Math. Ann. **145** 1961/1962, 429–439.
- [Br] Andreas Bredthauer, *Generalized Hyperkähler Geometry and Supersymmetry*, 15 pages, hep-th/0608114
- [D] Demailly, J.-P., *Regularization of closed positive currents and intersection theory*, J. Algebraic Geometry, **1** (1992), 361–409.
- [DP] Demailly, J.-P., Paun, M., *Numerical characterization of the Kähler cone of a compact Kähler manifold*, math.AG/0105176 also in Annals of Mathematics, **159** (2004), 1247–1274.
- [Do] A. Douady, *Le problème des modules pour les sous-espaces analytiques d'un espace analytique donné*, Ann. Inst. Fourier 16-1 (1966), 1–95.
- [DO] S. Dragomir, L. Ornea, *Locally conformal Kähler geometry*, Progress in Mathematics, 155. Birkhäuser, Boston, MA, 1998.
- [F1] A. Fujiki, *On Automorphism Groups of Compact Kähler Manifolds*, Inv. Math. 44 (1978) 225–258

- [F2] A. Fujiki, *Closedness of the Douady spaces of compact Kahler spaces*, Publ. Res. Inst. Math. Sci. **14** (1978/79), no. 1, 1–52.
- [F3] A. Fujiki, *On the Douady space of a compact complex space in the category C* , Nagoya Math. J. **85** (1982), 189–211
- [GHR] Gates, S. J., Jr.; Hull, C. M.; Roček, M., *Twisted multiplets and new supersymmetric nonlinear σ -models*, Nuclear Phys. B 248 (1984), no. 1, 157–186.
- [GP] Grantcharov, G., Poon, Y. S., *Geometry of hyper-Kähler connections with torsion*, math.DG/9908015, also in Comm. Math. Phys. 213 (2000), no. 1, 19–37.
- [GR] H. Grauert, R. Remmert, *Über kompakte homogene komplexe Mannigfaltigkeiten*, Arch. Math., 13 (1962) pp. 498–507
- [Hi] Hitchin, N. *Instantons, Poisson structures and generalized Kähler geometry* Comm. Math. Phys. 265 (2006), no. 1, 131–164, also in math.DG/0503432.
- [Ho] H. Hopf, *Über die Topologie der Gruppen-Mannigfaltigkeiten und ihrer Verallgemeinerungen*, Ann. of Math. 42 (1941), 22–52. Reprinted in Selecta Heinz Hopf, pp. 119–151, Springer, Berlin (1964).
- [Hö] Höfer, Thomas, *Remarks on torus principal bundles*, J. Math. Kyoto Univ. **33** (1993), no. 1, 227–259.
- [HP] Howe, P. S. Papadopoulos, G., *Twistor spaces for hyper-Kähler manifolds with torsion*, hep-th/9602108, also in Phys. Lett. B 379 (1996), no. 1–4, 80–86.
- [J] Joyce, Dominic, *Compact hypercomplex and quaternionic manifolds*, J. Differential Geom. **35** (1992) no. 3, 743–761
- [MV] Ruxandra Moraru, Misha Verbitsky, *Stable bundles on hypercomplex surfaces*, math.DG/0611714, 17 pages
- [Mum] D. Mumford, *Abelian Varieties*, Oxford University Press, Oxford, 1970.
- [OP] Opfermann, A., Papadopoulos, G., *Homogeneous HKT and QKT manifolds*, math-ph/9807026.
- [OV2] L. Ornea and M. Verbitsky, *An immersion theorem for compact Vaisman manifolds*, Math. Ann. **332** (2005), no. 1, 121–143, see also math.AG/0306077
- [Sa] Samelson, H., *A class of complex-analytic manifolds*, Portugal. Math. **12** (1953) 129–132.
- [SSTV] Ph. Spindel, A. Sevrin, W. Troost, A. Van Proeyen *Extended supersymmetric σ -models on group manifolds*, Nucl. Phys. B308 (1988) 662–698.
- [T] Tits, J. *Espaces homogènes complexes compacts*, Comment. Math. Helv. **37** 1962/1963, 111–120.

- [V] M. Verbitsky, *Stable bundles on positive principal elliptic fibrations*, 17 pages, math.AG/0403430, also in Math. Res. Lett. 12 (2005), no. 2-3, 251–264.
- [Wa] H. C. Wang, *Closed manifold with homogeneous complex structure*, Amer. J. Math., **76** (1954), 1-32.
- [Wo] Wolf, J. A., *On the geometry and classification of absolute parallelisms. I, II*, J. Diff. Geom. 6 (1971/72), 317-342, **7** (1972), 19-44.

INSTITUTE OF THEORETICAL AND EXPERIMENTAL PHYSICS,
B. CHEREMUSHKINSKAYA, 25, MOSCOW, 117259, RUSSIA
verbit@maths.gla.ac.uk, verbit@mccme.ru